

# RC RESEARCI \_\_\_\_\_

Bulletin #FO 086



# AVAILABLE WORKDAYS IN MISSISSIPPI FORESTS RESULTING FROM CLIMATE AND SOIL TYPE INTERACTIONS

Jonathan W. Pote Professor, Department of Agricultural and Biological Engineering and Director, Mississippi Water Resources Research Institute

> Charles L. Wax Professor and Head, Department of Geosciences and State Climatologist for Mississippi

Bob L. Karr Professor, Department of Forestry and Assistant Dean, College of Forest Resources

# Table of Contents

	Page
List of Figures	3
List of Tables	4
Abstract	5
Introduction	6
Background Information. The water balance technique Climate anti soil conditions	7 .8
Methods and Procedures	8
Model selection and assumptions Site selection Climatological data Model verification. Simulation Analyses of simulation	
Results and Discussion	<u>1</u> 3
Annual analyses Monthly analyses Applications of the information	13 16 18
Conclusions	1.8
References Cited	19

# List of Figures

Figure	Page
1. Location of the study sites	11
2. Minimum forest workdays annually, three probability levels, three soil types, three sites .	15
3. Minimum forest workdays monthly, three probability levels, three soil types, three sites.	17

# List of Tables

Tał	Page Page
1.	Comparison of observed and calculated soil moisture, selected dates
2.	Probability of minimum number of forest workdays, northern Mississippi, three soil types
3.	Probability of minimum number of forest workdays, central Mississippi, three soil types
4.	Probability of minimum number of forest workdays, southern Mississippi, three soil types14

#### Abstract

A computer model used to predict soil moisture was applied to three regions in Mississippi (northern, central, and coastal) and three soil textures (clay, loam, and sand). The model's performance was verified based on data from a previous study on forested lands in Mississippi. A day was considered suitable for harvest work if soil moisture was at or below 95% for sand, 80% for loam, and 65% for clay soils. A thirty year daily simulation (1961-90) provided calculated workdays for the nine combinations. These data were analyzed annually and monthly to produce the number of workdays expected at several probability levels.

Results indicate that sandy soils in the northern region allow 233 workdays/yr at the 50% probability level, whereas clay soils there allow only 140 days at that level. In the southern region sandy soils allow 252 workdays/yr at the 50% level and clay soils allow 122 days at that level. These results were produced at 10 probability levels for all soil types and all locations, as well as for each month.

#### Introduction

The warm, humid climate of the southern region of the U.S. is favorable to the growth of forests. The value of the timber produced annually is increasing and the region has recently surpassed the Northwest to lead the nation in lumber production (Forest and Wildlife Research Center, 1996). In the state of Mississippi, the net value added of the timber harvest reached an all-time high of \$4.6 billion in 1992, and more than seven million cords of pulpwood were produced that year (Barnett and Reinschmiedt, 1996).

Weather is the primary control affecting the timeliness of timber and pulpwood harvest. Weather exerts its control on harvest operations primarily through the soil moisture regime. Field work when the soil is too wet, even if physically possible, results in serious damage to the soil structure and increased time required for an operation (Hassan and Broughton, 1975). Soils that are too wet to support heavy equipment will be compressed and rutted during harvesting of trees. These physical alterations of soil lead to increased bulk density, loss of soil macroporosity, increased erosion, decreased availability of water, and potential loss of productivity (Rachael and Karr, 1989; Guo, et al, 1991; Karr, et al, 1991). Therefore, the obvious advantage of the forest productivity stemming from the humid climate type is often offset by the excess rainfall resulting in soils that are too wet for harvest operations.

The interaction between evapotranspiration, rainfall, and soil moisture storage influences tractability of soil and determines the number of workdays for a specific soil, providing periods of time on both annual and monthly scales when harvest is limited or possible. This water balance aspect of the natural environment is governed by climate, and the temporal pattern of occurrence should become evident by analysis of the corresponding climate record.

The availability of such information-probability of number of workdays on soil types in different locations-could allow Mississippi producers to make more informed decisions in several areas. For example, equipment purchases could be amortized according to the probable number of workdays and consequently the likely potential income from harvest operations. This advantage would benefit both the borrower and the lending institution. Also, the harvesting operator, mill owners, and other potential users of the forest products could use this type information to plan for more evenly distributed availability of materials through the entire year.

Therefore, the objective of this study is to establish the relationship that exists between climate and the pattern of workdays in the state, and then to provide guidance on the number of potential workdays based on sets of probability levels. This information will be organized by soil type and location, both monthly and annually.

#### Background Information

#### The water balance technique-

Water balance analyses evaluate the amounts of water added to landscapes by precipitation, stored in the environment as soil moisture, or lost from landscapes by evapotranspiration or runoff. Carried out daily, this assessment provides an accounting of moisture fluctuations resulting from climatic processes that determine the workability of soils for forest harvest. Data needed for water balance analyses include precipitation and energy inputs which are used to estimate evapotranspiration. Soil storage capacity is also needed to evaluate the condition of the soil and the resulting pattern of soil moisture for supporting harvest activities.

Studies of the water balance aspects of climate have been conducted for a variety of environmental and economic objectives such as reforestation on water yield from watersheds (Muller, 1966), groundwater availability (Robertson, 1973), streamflow analysis (Shelton, 1974), insect outbreaks (Kalkstein, 1974), and flood origins and analyses (Muller, 1976; Wax and Tingle, 1981). Growth in the use of water balance techniques has also led to application-oriented publications. For example, Muller and Larimore (1975) produced an atlas of seasonal water budget components of Louisiana, and Wax (1982) produced a similar document for Mississippi.

Recent use of the technique has been directed toward the immediate needs of industries and agriculture. The water balance approach has been used to assess the impact of climatic variability on aquacultural ponds seasonally, annually, and spatially (Pote and Wax, 1988; Wax and Pote, 1990). The results of this method have also been used to establish the drying potential, field work potential, and wet-dry periods that impact agricultural and forestry production on a daily basis (Wax, et al, 1990). Irrigation system design criteria have been established using water balance methods (Pote and Wax, 1986). The method has been used to investigate opportunity for land application of wastewater (Pote and Wax, 1995) and to model quantity and quality of overflow from aquacultural ponds (Pote, et al, 1996). The water balance method of climatic analysis, therefore, is a useful tool for investigating atmospheric interactions with environmental processes and economic endeavors of humans.

#### Climate and soil conditions--

In Mississippi a strong gradient of precipitation and evaporation occurs from the Gulf Coast to the interior that strongly influences the water balance patterns across the state (Wax and Pote, 1996). Soils in the state are highly variable in texture both locally and regionally. The resulting marked differences in permeabilities cause large variations in the soil moisture storage component of the water balance. Therefore, moisture, energy, and soil moisture conditions are often found in many combinations which affect tractability and consequently the number of field working days.

The Canadian Society of Soil Science (1967) considers soil tractable if a tractor or other forest machine can move on that soil to satisfactorily perform the function of the machine without causing significant damage to the soil. Elliot, et al (1977) suggested that the weather has a considerable effect on available field working days, particularly on certain soil types during months with high precipitation. Rutledge and McHardy (1968) indicated that the soil is not tractable when soil moisture is near or above field capacity. Subsequently they defined the criterion of a workday as one in which the moisture content was less than 95% of capacity. Bolton, et al (1968) classified a day as a workday if soil moisture in the top six-inch layer was less than 80% of field maximum for

tillage operations on silty clay and sandy soils. For clay soils, the moisture must be at or below 78% of field maximum for all field operations.

Knowing the probable number of forest workdays at a given location can have far-reaching economic effects on a harvest operation. These include decisions on amortization of equipment or on obtaining loans for equipment, harvest scheduling by location and soil type, value of leases, amount of stockpiling required in woodyards, and other financial concerns facing Mississippi owner/operators.

#### Methods and Procedures

#### Model selection and assumptions-

Matsumoto (1992) first attempted an analysis of available forest working days by using sites and conditions in Mississippi. That study used an original water balance model and assumed a 95% moisture level criterion for all soil types to model daily water balance components over the 20-year period 1970-1989. For this study it was deemed important to use a well-established and flexible model which would allow setting parameters to better account for complex conditions of climate and soils across the state, and to expand the period of analysis to a standard 30-year interval.

McCabe, et al, (1985) published a continuous daily water budget model through the Southern Regional Climate Center which uses the Thornthwaite method (Thornthwaite and Mather, 1957) of computing potential evapotranspiration. That model was chosen for this study because it allows the capability of setting multiple parameters, therefore providing maximum accounting of the varying soil and climate conditions in the state. The model requires inputs of only daily precipitation and maximum and minimum temperatures, and allows the user to set soil moisture storage conditions such as soil depth and water holding capacity. It also provides the flexibility of setting two layers of soil moisture, the upper with equal availability of soil moisture (available on-demand), and the lower with soil moisture becoming decreasingly available.

The following assumptions were made for this study:

- Soil moisture storage capacity was set at 3" (1.2 inches per foot throughout an effective root zone of 2.5 feet) for sandy soil, 6" for loamy soil (2.4 inches per foot, effective rooting zone of 2.5 feet), and 9" for clay soil (3.6 inches per foot, effective rooting zone of 2.5 feet);
- the upper layer of soil moisture storage capacity was set at I" for all texture classes, with the balance assigned to the lower level;
- 3) the tractability threshold was set for sandy soil at 95% of the capacity (2.85"), for loamy soil at 80% of the capacity (4.8"), and for clay soil at 65% of the capacity (5.85")-these levels are slightly more conservative for clay soils than those in Matsumoto (1992) and Bolton, et al (1968);
- soil moisture storage in all soils was at capacity on the first day of the period of each simulation;
- 5) point locations were used to represent the zones of the state;
- 6) no accounting was made for the effect of slope in the analyses; and
- 7) all workdays are of equal length.

#### Site selection-

Sites were chosen to represent climate conditions across Mississippi. Criteria for site selection were: **1**) spatial dispersion providing representation of the climatic gradient from the coast to the interior; and, 2) availability of serially complete and homogeneous daily climatological data in digital format.

Based on the best possible combination of these criteria, the following sites in Mississippi were selected: Poplarville representing the coastal zone, State University representing the interior zone, and Meridian between the two (figure 1). All three soil textures are found in forested areas in the vicinity of each of these locations.

#### Climatological data-

Daily precipitation and maximum and minimum temperatures for the period 1961-1990 were obtained from observations of the National Weather Service Cooperative Observation System stored on CD-ROM (Earth info, 1992). There were cases where data were missing for up to a month. In these cases, data from the next-nearest location on the same dates were substituted to make the records serially complete. It is estimated that less than 0.5% of the data were thus derived.

#### Model verification-

In her study, Matsumoto (1992) measured soil moisture on 10 different days in 1992 at three soil texture sites for comparison with her calculated data for the same times and soil types. For purposes of verification, the model adopted for this present study was run three times for the entire year 1992, with soil moisture capacity set for 3, 6, and 9 inches, respectively. Calculated soil moisture was then extracted for each of those 10 sampling days for comparison with the measured data. Observed and calculated results matched on 29 of the 30 sample dates, providing a high level of confidence that the model was accurately simulating the pattern of soil moisture in the forest locations.

In field sampling, Matsumoto (1992) used a moisture level greater than 95% of field capacity as the criterion on all soils to designate an unsuitable workday; less than 95% indicated a suitable workday. The criteria used in this study to designate an unsuitable workday were as follows: soil moisture greater than 95% for sandy soils, 80% for loamy soils, and 65% for clay soils. Table 1 shows a comparison of Matsumoto's measured results along with the results calculated by the model used in this study.



	<u>Sandy</u>		Loamy		Clav	
<u>Date</u>	<u>Obs</u>	Model	<u>Obs</u>	Model	<u>Obs</u> N	1 <u>odel</u>
2/6	U	U	U	U	U	U
2/10	S	U	U	U	U	U
2/14	U	U	U	U	U	U
2/19	U	U	U	U	U	U
2/24	U	U	U	U	U	U
2/26	U	U	U	U	U	U
3/11	U	U	U	U	U	U
4/8	U	U	U	U	U	U
4/10	S	S	U	U	U	U
4/15	S	S	S	S	S	S

Table 1: Comparison of Observed and Calculated Soil Moisture, Selected Dates

U = Conditions unsuitable for a work day

S = Conditions suitable for a work day

#### Simulation-

The model simulated the water balance components daily over the 30-year period **1961**-1990. For each of the three locations, the 30-year period was modeled multiple times to estimate soil moisture in sandy (3" capacity), loamy (6" capacity), and clay (9" capacity) soils. The combination of three soil texture classes and three locations required that the model be run nine times to produce the needed data.

#### Analyses of simulation-

Daily soil moisture storage estimates were separated from the remainder of the water balance components and put into electronic spreadsheets, one for each of the three soils at each of the three sites. Each day's soil moisture value in these nine spreadsheets was then subjected to the criteria for suitable or unsuitable workdays. These results were counted and tabulated, first annually and by months, resulting in 30 numbers for annual values and 30 values for each month. These 30 yearly and monthly data were ranked in descending order, then divided into 10% quantiles from which empirical probabilities could be determined. Probability tables and graphical analyses were thus produced to show potential for workdays on all three soil types, both spatially and temporally.

#### Results and Discussion

#### Annual analyses-

Results of the annual analysis for northern Mississippi are shown in Table 2. The study shows that not only are there fewer suitable working days on clay soils, but also that the number decreases dramatically at the higher probability levels. For example, at least 192 days are expected each year at the 10% level on clay soils, while only 52 are expected at the 90% level. In comparison, sandy soils in that zone show much less variability in number of annual working days at the different probabilities. At least 260 are expected 10% of the time on sandy soils and at least 216 are expected at the 90% level. On loamy soils the equivalent values are 233 and 141, respectively.

Probability of Minimum Types	Number of Forest	Workdays, Northern	i Mississippi, Three Soil
% Chance	Sandy	Loamy	Clay
10	260	233	192
20	256	224	179
30	244	201	161
40	240	193	149
50	233	183	140
60	230	177	137
70	223	167	123
80	217	157	99
90	216	141	52
	Probability of Minimum Types % Chance 10 20 30 40 50 60 70 80 90	Probability of Minimum Number of Forest Types   % Chance Sandy   10 260   20 256   30 244   40 240   50 233   60 230   70 223   80 217   90 216	Probability of Minimum Number of Forest Workdays, Northern Types   % Chance Sandy Loamy   10 260 233   20 256 224   30 244 201   40 240 193   50 233 183   60 230 177   70 223 167   80 217 157   90 216 141

Table 3 shows results for central Mississippi. On clay soils the variability of number of workdays is even more dramatic. At the 90% probability level at least 200 workdays/year are expected, while at the 10% level only 7 workdays/year are anticipated. On sandy soils in that zone, the expected workdays/year are 282 and 225, respectively, at the two probability levels. Loamy soils have equivalent values of 239 and 108 for those probability levels.

% Chance	Sandy	Loamy	Clay	
10	273	238	221	
20	260	221	191	
30	253	205	176	
40	250	197	147	
50	243	192	142	
60	232	179	135	
70	228	163	122	
80	227	154	99	
90	219	143	88	

Table 3: Probability of Minimum Number of Forest Workdays, Central Mississippi, Three Soil Types

In southern Mississippi (Table 4) clay soils allow 221 and 88 workdays/year at the 90% and 10% levels of probability. On sandy soils harvesters can expect 273 and 219 workdays/year at these two levels. Loamy soils provide at least 219 and 273 workdays/year at these levels of probability in this zone.

Table 4:	Probability of Minimum Number of Forest Workdays, Southern Mississippi, Three Soil Types					
	% Chance	Sandy	Loamy	Clay		
	10	282	239	200		
	20	274	209	175		
	30	267	200	147		
	40	254	180	134		
	50	252	172	122		
	60	238	166	98		
	70	234	140	82		
	80	232	120	43		
	90	225	108	7		

2

The number of workdays expected annually in the three zones at the 80%, 50%, and 20% probability levels is shown in Figure 2 for comparison. These results indicate that clay soils in all zones are limiting, and harvesters with holdings on these soils will need to be especially careful to take advantage of the windows of opportunity that are available through the year. Sandy and loamy soils, on the other hand, offer so many more potential workdays that balancing holdings in mixed soil types, if possible, would strongly enhance harvest operations throughout the state.



Even though there are, on the average, limitations on work days on all soil types and particularly on clay soils, results indicate that over a number of years there are a remarkably high number of workdays available on all soils, even on clay soils or in the worst of cases. While any given or single year may not fulfill such positive expectations, with the right timing or planning there is systematically better access to all soil types at all locations than may be generally recognized.

Of particular interest, it can be seen that sandy soils allow more potential workdays/year in the southern portion of the state than in the northern portion of the state while clay soils allow more workdays/year in the north than in the south (Figure 2). The most likely explanation for this phenomenon is climatic in nature. Higher annual rainfall amounts in the coastal region keep clay soils near saturation a higher percentage of the time there, whereas higher average temperatures cause sandy soils to dry more rapidly in that region on an annual basis. Since one season may be dominant enough to skew the annual averages, these phenomena may be better understood by comparison of seasonal or monthly results rather than annual results.

#### Monthly analyses-

Figure 3 shows monthly probabilities of workdays at three levels for three soils in all three zones of the state. It is immediately obvious that sandy soils exhibit consistently high probability for workdays throughout the state with few limitations from about mid-March through November. Also, the sandy soils exhibit very little difference at all three levels of probability. By comparison, the high probability period on loamy soils is reduced by over a month in the spring. Additionally, moving from north to south there is a noticeable loss of probable workdays at the 80% level compared to the other levels. An even stronger contrast is seen on clay soils, as the high probability period is reduced by yet another month in the spring throughout the state and the 80% level practically disappears in the coastal zone of the state.

This means that harvesting in northern Mississippi is limited very little by soil type at the 20% and 50% levels, with more limitations imposed at the 80% level. Also, sandy soils allow considerably earlier access in that portion of the state. In contrast, sandy soils in southern Mississippi offer more workdays in the spring at lower probability levels, but working conditions on clay soils may severely limit operations at the highest probability levels almost any time during the year. Therefore, operators in the coastal zone have an advantage on sandy soils throughout the year, while operators in the north have a much more limited season of workdays in general but have the advantage of a high probability of being able to harvest on clay soils throughout the year.

#### Applications of the information-

For the independent owner-operator, this information should be useful in predicting how to plan loan repayments. Rather than hoping for a consistently high level of workdays, there will be some sense of the level of risk in this assumption. if the operators have access to a variety of soil types, the information can also be used to aid in harvest scheduling to stabilize income.

This information obviously holds similar value for the lending agency. For example, the risk associated with various amortization schedules can be quantified at selected probability levels. The potential for return on investment can more easily be assessed using these illustrated levels of risk.

On clay soils, the lending agency is likely to prefer predictions based on conservative estimates (high probability, low risk situations). In reality the harvester can systematically enhance



chances for profit by correctly taking advantage of the windows of opportunity presented by climatic variability in the region.

For the landowner, these data help account for the variation in value of land. Knowing that operators will be saving sandy soils for certain times of year will increase its value at those times. In addition, size of equipment and the time required for harvest as a consequence, will produce a trade off between land value and equipment type.

The owner of a sawmill or a chipping operation can use the information to make determinations at different risk levels about how much raw material should be stockpiled in their yards during various seasons. This capability should help to stabilize employment and income in this segment of the industry.

#### Conclusions

The following conclusions are drawn based on the data and analyses:

- 1) This model can be used to predict forest workdays at different levels of probability on different soils across Mississippi.
- 2) The predicted values are considered valid because the analyses are based not on a theoretical distribution but on actual daily climatic records over a 30-year period.
- 3) Overall, the analyses show the expected trend of sandy soils providing more probable workdays than clay soils. Unexpectedly, the analyses showed an increase in probable workdays on sandy soils going south across the state, whereas the number of workdays on clay soils increased going north. This finding is considered to be climatological in origin.
- 4) At lower probability levels (20%, 50%), few unexpected findings resulted either by soil type or by location in the state. In contrast, at high probability levels (80%) dramatic changes in workday potential were evident by both soil type and location.
- 5) Trends resulting from the annual analyses were dominated by seasonal or monthly variation. Specifically, the results showed loss of virtually all workdays on clay soils in winter over the entire state, with a similar yet less pronounced pattern on loamy soils. Furthermore, workdays on clay soils at the 80% probability level were highly variable from north to south in summer. For example, in August expected workdays vary from almost a full month in the north and central zones to near zero in the southern portion of the state.
- 6) These results show very positively that there are more opportunities for harvest operations than are generally realized even on the most limiting soils. Therefore it is concluded that if these opportunities are systematically maximized, significantly increased or stabilized incomes could potentially result.
- 7) Overall the results of this study appear to offer significant information to several aspects of the industry. These include land owners, timber harvesters, sawmill and chipping mill owners, and lenders, all of whom have different perspectives on the number of potential workdays in the forests.

#### **References** Cited

Barnett, B. and L. Reinschmiedt. 1996. Agriculture in Mississippi, an Analysis of the Impacts of Food and Fiber Related Sectors on the Mississippi Economy. Department of Agricultural Economics, Mississippi State University, Mississippi State, MS.

Bolton, B., J.B. Penn, F.T. Cooke, Jr., and A.M. Heagler. 1968. Days suitable for fieldwork-Mississippi River delta cotton area. *Dep. Agri. Econ. and Agribusiness*. Louisiana State University, Baton Rouge, LA. D. A. E. Research Report #384.

Canadian Society of Soil Science. 1967. *Glossary of soil terms in Canada*. Prepared by the Glossary subcommittee.

Earth Info. 1992. NCDC Summary of the Day CD-ROM. Earth Info, inc., Boulder, CO.

Elliot, R.L., W.D. Lembke and D.R. Hunt. 1977. A simulation model for predicting available days for soil tillage. Trans. ASAE, 20:4-8.

Ferguson, Bruce K. 7 996. Estimation of direct runoff in the Thornthwaite water balance. Professional *Geographer*, 48(3): 263-271.

Forest and Wildlife Research Center. 1996. *Discover* Mississippi's *Forests.* Forest and Wildlife Research Center, Mississippi State University, Mississippi State, MS.

Guo, Y., B.L. Karr and J.M. Rachael. 1991. Effects of soil compaction and the presence or absence of the A soil horizon on water relations of Loblolly Pine seedlings in North-Central Mississippi. In S.E. Coleman and D.G. Neary (eds.), *Proceedings* of the *6th* Biennial Southern *Silvicultural* Research Conference. Memphis, TN, pp. 449-457.

Hassan, A.E. and R.S. Broughton. 1975. Soil moisture criteria for tractability. Can. *Agric. Eng.* 17(2):124-1 29.

Kalkstein, L.S. 1974. The effect of climate upon outbreaks of the southern pine beetle. Publications in Climatology, 27(3). Laboratory of Climatology, Centerton, NJ.

Karr, B.L., J.M. Rachael and Y Guo. 1991. Modeling bulk density, macroporosity, and microporosity with the depth of skidder ruts, for a loess soil in North Central Mississippi. In S.E. Coleman and D.G. Neary (eds.), *Proceedings* of the *6th Biennial* Southern *Silvicultural Research Conference*. Memphis, TN, pp. 443-448.

Matsumoto, S. 1992. Estimation of suitable field working days for timber harvest. Masters thesis, Mississippi State University, Mississippi State, MS.

McCabe, G.J., Jr., J.D. McLaughlin and R.A. Muller. 1985. Thornthwaite continuous daily water budget: A computer program for microcomputers. Louisiana Office of State Climatology, Miscellaneous Publication: May 1985. Louisiana State University, Baton Rouge, LA.

Muller, R.A. 1966. The effects of reforestation on water yield: A case study using energy and water balance models for the Allegheny Plateau, New York. Publications in *Climatology*, 19(3). Laboratory of Climatology, Centerton, NJ.

Muller, R.A. 1976. Comparative climatic analysis of lower Mississippi River floods: 1927, 1973, and 1975. Water Resources Bulletin, 12(6). American Water Resources Association.

Muller, R.A. and P.B. Larimore. 1975. Atlas of seasonal water budget components of Louisiana. (1941-1970). Publications in Climatology, 28(1). Laboratory of Climatology, Centerton, NJ.

Pote, J.W. and C.L. Wax. 1986. Climatological aspects of irrigation design criteria in Mississippi. Technical Bulletin 138. Miss. Ag. and Forestry Experiment Station, Mississippi State University, Mississippi State, MS.

Pote, J.W. and C.L. Wax. 1995. Climatic Criteria for Land Application of Municipal Wastewater Effluent. *Water Research*, Vol 29, No.1, pp. 323-328.

Pote, J.W., C.L. Wax and C.R. Tucker. 1988. Water in Catfish Production: Sources, Uses, and Conservation. Special Bulletin 88-3, Miss. Ag. and Forestry Experiment Station, MSU, Miss. State, MS.

Rachael, J.M. dnd B.L. Karr. 1 989. Effects of current harvesting practices on the physical properties of a Loessal soil in west-central Mississippi. In James E. Miller (ed.), Proceedings of the 5th Biennial Southern Silvicultural Research Conference. Memphis, TN, pp. 527-532.

Robertson, D.K. 1973. Groundwater availability in southern New Jersey. *Publications in climatology*, 26(1), Laboratory of Climatology, Centerton, NJ.

Ruthedge, P.L. and F.V. McHardy. 1968. The influence of the weather on field tractability in Alberta. Can. Agric. Eng., 10(2): 70-73.

Thornthwaite, C.W. and J.R. Mather. 19.57. Instructions and tables for computing potential evapotranspiration and the water balance. *Publications* in *Climatology*, 1 0(3), Laboratory of Climatology, Centerton, NJ.

Tucker, C.S., S.K. Kingsbury, J.W. Pote and C.L. Wax. 1996. Effluents from Cannel Catfish Ponds in Northwest Mississippi. *Aquaculture*.

Wax, Charles L. 1982. Atlas of the Climatic Water Balance in Mississippi: Normals and Variability, 1951-1980. Water Resources Research institute Technical Publication DO 18. Mississippi State University, Mississippi State, MS, 82 pp.

Wax, C.L. and J.L. Tingle. 1981. The Pearl River flood of Easter, 1979: A water balance explanation. Water Resources Research Institute Bulletin, Mississippi State University, Mississippi State, MS.

Wax, C.L. and J.W. Pote. 1990. A Climatological Basis for Conservation of Groundwater in Aquacultural Production in the Southern Region." Technical Bulletin 169 Miss. Ag. and For. Experiment Station, MSU, Miss. State, MS.

Wax, C.L. and J.W. Pote. 1996. The influence of Climate on Design of Systems for I-and Application of Wastewater. *Climate Research*, 6:71-78.

Wax, C.L., J.W. Pote and R.A. Poncet. 1 990. Selected Climatological Features Affecting Agriculture and Forestry in Mississippi. Bulletin 969. Miss. Ag. and For. Experiment Station, MSU, Miss. State, MS.

![](_page_22_Picture_0.jpeg)

Forest and Wildlife Research Center Mississippi State University Box 9680 Mississippi State, Mississippi 39762

# Mississippi State

Mississippi State University does not discriminate on the basis of race, color, religion, national origin, sex, age, disability, or veteran status

72768/9-97